

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total
Injury and Violence

Health Risk Behavior and Percentages

Linear Change*

Quadratic Change*

Change from
2017-2019 †

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)

23.5 18.7 18.1 17.6 13.4 12.4 10.1 9.1 8.7 7.8 Decreased, 1997-2019 No quadratic change No change

QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)

5.7 5.9 3.9 4.2 No linear change Not available[§] No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]												
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019															
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)															36.3	36.5	34.7	32.2	Decreased, 2013-2019			Not available [§]			No change				
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)															26.4	18.5	23.1	24.4	21.7	22.8	20.7	23.1	20.5	15.7	Decreased, 1997-2019			No quadratic change	Decreased
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)															15.2	7.4	6.8	8.0	6.5	7.4	6.4	6.5	4.9	2.7	Decreased, 1997-2019			No quadratic change	Decreased
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)															7.3	5.3	No linear change			Not available			No change						

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)																	
			6.8			5.2	8.0	8.3	7.9	7.4	5.4	7.2	7.1	7.1	No linear change	No quadratic change	No change
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)																	
			32.5			26.4	29.6	27.0	28.7	28.7	21.2	19.9	21.4	19.7	Decreased, 1997-2019	No quadratic change	No change
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)																	
			12.5			10.1	12.7	10.6	9.5	11.4	6.0	7.8	7.7	5.6	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total
Injury and Violence

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)																	
													10.0	8.6	No linear change	Not available [§]	No change
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											9.8	10.1	6.5	6.8	Decreased, 2013-2019	Not available	No change
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											9.8	8.9	8.6	7.4	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)																	
								20.8	18.9	21.4	22.0	21.2	23.8		Increased, 2009-2019	No quadratic change	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)																	
										17.4	13.2	17.0	18.2	18.0	Increased, 2011-2019	Not available [§]	No change
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
					30.1	28.2	29.4	26.7	27.0	25.7	31.3	29.2	37.2		Increased, 2003-2019	Decreased, 2003-2013 Increased, 2013-2019	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
			22.4			17.6	15.4	15.1	14.6	14.8	15.0	15.7	14.8	18.4	Decreased, 1997-2019	Decreased, 1997-2007 No change, 2007-2019	Increased
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
			17.0			14.5	11.6	11.9	12.5	13.7	12.6	13.9	13.0	15.9	No linear change	Decreased, 1997-2007 Increased, 2007-2019	Increased
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)																	
			8.0			10.3	9.2	7.6	8.8	10.9	7.7	9.4	7.9	8.1	No linear change	No quadratic change	No change
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)																	
			2.5			3.7	2.7	2.4	3.5	4.6	2.9	3.9	2.8	3.0	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total Tobacco Use

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
		77.3				71.1	59.5	62.2	59.0	59.2	47.1	44.1	40.5	30.6	Decreased, 1997-2019	Decreased, 1997-2011 Decreased, 2011-2019	Decreased
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													15.5	11.8	Decreased, 2017-2019	Not available [§]	Decreased
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
		47.0				32.7	26.2	26.0	26.1	24.1	17.9	16.9	14.3	8.9	Decreased, 1997-2019	No quadratic change*	Decreased
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
		27.6				18.4	14.4	13.4	12.0	11.6	7.3	5.7	4.9	3.0	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)																	
			22.5			14.9	11.1	10.6	9.3	9.0	5.6	4.6	3.7	2.2	Decreased, 1997-2019	No quadratic change	No change
QN33: Percentage of students who smoked more than 10 cigarettes per day (on the days they smoked during the 30 days before the survey, among students who currently smoked cigarettes)																	
			26.2			16.6	19.1	13.6	13.8	18.2	10.9	6.8	7.2	11.0	Decreased, 1997-2019	No quadratic change	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])																	
												41.7	44.5	53.7	Increased, 2015-2019	Not available [§]	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)															23.4	14.1	26.1	No linear change			Not available [§]			Increased
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)															3.2	2.7	11.1	Increased, 2015-2019			Not available			Increased
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)															2.2	1.9	8.7	Increased, 2015-2019			Not available			Increased
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)															29.4	21.0	28.1	No linear change			Not available			Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

†Based on t-test analysis, $p < 0.05$.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Tobacco Use**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019											
QN36: Percentage of students who usually got their own electronic vapor products by buying them in a store (such as a convenience store, supermarket, discount store, gas station, or vape store, during the 30 days before the survey, among students who currently used electronic vapor products and who were aged <18 years)														14.0	12.3	No linear change	Not available [§]	No change							
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)														10.6	6.4	Decreased, 2017-2019	Not available	Decreased							
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)														18.7	15.5	15.5	17.2	17.5	13.3	14.0	11.0	7.9	Decreased, 2003-2019	No change, 2003-2011 Decreased, 2011-2019	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)																	
						39.1	30.8	31.3	31.5	30.3	23.9	23.4	18.2	13.0	Decreased, 2003-2019	Decreased, 2003-2015 Decreased, 2015-2019	Decreased
QN39: Percentage of students who tried to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)																	
												45.8	50.5		No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total Alcohol and Other Drug Use																Linear Change*	Quadratic Change*	Change from 2017-2019[†]
Health Risk Behavior and Percentages																		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																		
			30.4			26.8	28.9	25.1	21.7	20.5	19.0	15.9	17.1	15.4	Decreased, 1997-2019	No quadratic change	No change	
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)																		
			49.3			45.1	37.4	40.6	37.8	34.6	30.4	28.5	26.6	23.5	Decreased, 1997-2019	No quadratic change	No change	
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																		
													13.2	10.3	Decreased, 2017-2019	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total Alcohol and Other Drug Use															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN45: Percentage of students who ever used marijuana (one or more times during their life)																	
			48.1			43.3	34.4	35.0	31.4	37.4	34.0	33.1	32.1	31.9	Decreased, 1997-2019	Decreased, 1997-2007 Decreased, 2007-2019	No change
QN46: Percentage of students who tried marijuana for the first time before age 13 years																	
			9.3			11.5	10.0	10.2	9.2	10.0	8.5	6.3	7.4	7.0	Decreased, 1997-2019	No change, 1997-2004 Decreased, 2004-2019	No change
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)																	
			28.6			21.1	15.8	16.4	16.1	19.2	17.7	17.2	15.8	16.1	Decreased, 1997-2019	Decreased, 1997-2004 No change, 2004-2019	No change
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)																	
												10.3	7.9	6.2	Decreased, 2015-2019	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total Alcohol and Other Drug Use															Linear Change*	Quadratic Change*	Change from 2017-2019 †	
Health Risk Behavior and Percentages																		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)																		
													10.9	11.0	No linear change	Not available [§]	No change	
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)																		
		8.3				9.8	8.3	8.6	6.0	7.5	4.5	4.6	4.7	3.5	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change	
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)																		
		24.7				14.3	13.5	14.2	13.4	11.4	7.1	7.4	6.5	5.6	Decreased, 1997-2019	No quadratic change	No change	
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)																		
													3.7	2.1	1.8	Decreased, 2015-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total Alcohol and Other Drug Use																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
						6.7	5.9	6.5	6.9	6.6	4.2	5.0	4.2	3.6	Decreased, 2003-2019	No change, 2003-2009 Decreased, 2009-2019	No change
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
			6.1			7.1	5.7	6.1	4.0	5.3	2.9	4.5	3.6	2.0	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	Decreased
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													2.5	2.2	No linear change	Not available [§]	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
			34.2			30.4	19.8	27.0	25.6	24.4	20.6	20.9	22.4	24.0	Decreased, 1997-2019	Decreased, 1997-2015 No change, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total
Sexual Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN58: Percentage of students who ever had sexual intercourse																	
			53.7			52.1	46.3	50.3	48.3	51.8	44.7	41.7	38.4	39.2	Decreased, 1997-2019	No quadratic change	No change
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years																	
			7.2			5.6	7.9	7.8	6.7	7.2	5.4	4.8	3.7	3.2	Decreased, 1997-2019	No change, 1997-2009 Decreased, 2009-2019	No change
QN60: Percentage of students who had sexual intercourse with four or more persons during their life																	
			18.1			14.8	13.6	14.4	12.7	16.6	12.6	10.4	9.6	8.5	Decreased, 1997-2019	No quadratic change	No change
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)																	
			39.4			38.7	33.5	36.5	33.6	37.7	31.7	30.3	29.0	28.7	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total
Sexual Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)																		
			27.6			21.9	19.3	19.0	22.9	16.8	18.7	17.1	17.5	14.9	Decreased, 1997-2019	No quadratic change	No change	
QN63: Percentage of students who used a condom during last sexual intercourse (among students who were currently sexually active)																		
			57.3			61.7	65.2	59.0	59.9	50.6	53.1	53.9	48.7	52.6	Decreased, 1997-2019	Increased, 1997-2004 Decreased, 2004-2019	No change	
QN64: Percentage of students who used birth control pills before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																		
			18.5			17.5	18.4	20.5	23.4	21.2	19.9	22.4	24.5	24.6	Increased, 1997-2019	No quadratic change	No change	
QNIUDIMP: Percentage of students who used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																		
												2.6	5.2	7.7	6.6	Increased, 2013-2019	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Sexual Behaviors**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019									
QNSHPARG: Percentage of students who used a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active))														4.9	6.1	6.0	6.3	No linear change				Not available [§]	No change
QNOTHHPL: Percentage of students who used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)														27.3	33.6	38.3	37.5	Increased, 2013-2019				Not available	No change
QNDUALBC: Percentage of students who used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)														6.6	11.6	11.3	11.2	Increased, 2013-2019				Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Total
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)																	
			13.9			13.9	11.4	13.7	13.0	16.2	15.1	14.5	16.5	13.6	No linear change	No quadratic change	No change
QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)																	
												11.6	11.3	10.8	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						15.2	16.8	16.3	15.4	15.4	15.4	17.0	16.0	17.8	No linear change	No quadratic change	No change
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						14.5	15.4	15.4	17.4	16.5	18.0	18.5	20.2	18.4	Increased, 2003-2019	No quadratic change	No change
QN67: Percentage of students who described themselves as slightly or very overweight																	
			32.0			33.8	33.2	31.2	30.1	30.0	32.3	32.0	33.7	34.7	No linear change	No change, 1997-2011 Increased, 2011-2019	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)																	
						25.8	27.2	28.8	26.1	26.5	28.5	28.3	31.3	32.7	Increased, 2003-2019	No change, 2003-2015 Increased, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)																	
						18.5	20.8	18.9	18.8	19.1	14.2	14.8	16.7	18.4	Decreased, 2003-2019	No quadratic change	No change
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						9.9	11.3	10.6	9.7	9.7	8.0	7.7	8.8	10.4	Decreased, 2003-2019	No quadratic change	No change
QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						46.0	50.1	47.0	50.8	50.3	55.4	52.7	48.3	47.5	No linear change	Increased, 2003-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						19.2	23.8	21.3	24.2	23.0	25.7	21.0	20.8	20.9	No linear change	Increased, 2003-2013 Decreased, 2013-2019	No change
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)																	
						40.3	47.2	46.3	45.6	45.6	47.0	49.6	49.7	50.9	Increased, 2003-2019	No quadratic change	No change
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)																	
						25.6	24.2	24.3	23.1	25.7	26.5	29.0	35.4	37.6	Increased, 2003-2019	No change, 2003-2009 Increased, 2009-2019	No change
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)																	
						59.2	59.8	60.0	58.6	58.9	55.9	58.2	61.7	62.9	No linear change	No change, 2003-2015 Increased, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
						17.9	19.3	20.7	21.7	22.9	21.1	21.9	24.1	24.4	Increased, 2003-2019	No quadratic change	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						6.4	7.5	6.7	6.3	7.0	6.2	8.5	10.3	11.0	Increased, 2003-2019	No change, 2003-2013 Increased, 2013-2019	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						55.6	58.0	57.4	58.7	56.8	57.3	57.1	50.7	48.2	Decreased, 2003-2019	No change, 2003-2015 Decreased, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						22.2	26.3	22.0	22.3	25.4	22.7	23.2	18.3	18.7	Decreased, 2003-2019	No change, 2003-2015 Decreased, 2015-2019	No change
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						10.5	13.5	11.1	10.9	12.3	11.6	11.1	9.0	8.9	Decreased, 2003-2019	No change, 2003-2013 Decreased, 2013-2019	No change
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								16.4	18.1	17.6	21.7	23.6	23.2	25.4	Increased, 2007-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total															Linear Change*	Quadratic Change*	Change from 2017-2019[†]				
Weight Management and Dietary Behaviors																					
Health Risk Behavior and Percentages																					
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019							
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																					
						40.5		35.7		36.4		32.9		32.4		28.1		25.6	Decreased, 2007-2019	No quadratic change	No change
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																					
						31.4		26.7		26.3		24.5		23.3		20.5		17.6	Decreased, 2007-2019	No quadratic change	No change
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)																					
										21.9		22.5		26.0		30.0			Increased, 2013-2019	Not available [§]	Increased
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																					
										34.5		30.3		28.0		24.6			Decreased, 2013-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Total Weight Management and Dietary Behaviors															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNMILK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
									10.5	8.7	7.1	7.3	Decreased, 2013-2019		Not available [§]	No change	
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)																	
								14.9	15.5	12.3	12.6	15.4	16.6	No linear change		Decreased, 2009-2015 Increased, 2015-2019	No change
QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)																	
								31.9	29.9	40.1	34.8	30.2	26.5	Decreased, 2009-2019		Increased, 2009-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Physical Activity**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															39.3	39.8	37.0	40.6	37.4	No linear change			Not available [§]	No change
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															20.6	19.9	16.5	19.2	19.1	No linear change			Not available	No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															21.9	22.5	20.2	22.0	19.0	No linear change			Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Physical Activity**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																	
						30.8	35.5	27.4	28.8	32.0	26.7	25.5	20.9	21.3	Decreased, 2003-2019	No quadratic change	No change
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																	
								21.3	23.0	31.2	34.5	40.1	41.2	47.5	Increased, 2007-2019	No quadratic change	Increased
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																	
			31.3			34.9	25.2	31.0	32.9	35.4	34.4	37.8	31.2	31.5	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Total
Physical Activity

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)																	
			22.3			23.8	17.3	20.0	23.1	20.0	19.3	22.3	19.2	17.8	No linear change	No quadratic change	No change
QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)																	
						50.9	52.6	48.6	48.2	46.3	50.8	50.8	48.3	45.9	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Total Other	Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)																	
								69.7	69.0	67.8	69.4	69.8	73.1	71.6		Increased, 2007-2019	No quadratic change	No change
	QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)																	
								2.7	2.0	3.2	2.3	2.2	2.1	1.9		No linear change	No quadratic change	No change
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma																	
								22.1	26.1	24.3	26.7	23.1	25.5	26.0	23.9	No linear change	No quadratic change	No change
	QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)																	
													24.3	22.0	19.4	Decreased, 2015-2019	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Total Other	Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)																	
										67.3	73.1	73.8	76.8	78.2	Increased, 2011-2019	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Site-Added**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														59.5	62.7	No linear change	Not available [§]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														49.5	54.4	No linear change	Not available	No change
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														37.2	40.7	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

†Based on t-test analysis, $p < 0.05$.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Site-Added**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]										
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019													
QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)															20.8	17.2	17.6	No linear change	Not available [§]	No change							
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)															25.3	14.3	9.7	9.5	9.4	9.3	5.0	7.0	5.7	4.6	Decreased, 1997-2019	Decreased, 1997-2004 Decreased, 2004-2019	No change
QN93: Percentage of students who had oral sex															49.1	51.2	41.8	41.7	39.3	39.4	Decreased, 2009-2019	No quadratic change	No change				
QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)															9.2	9.0	No linear change	Not available	No change								

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Total
Site-Added**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]			
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																			
								76.7	77.3	74.5	77.2	72.9	70.6	70.4	Decreased, 2007-2019	No quadratic change	No change		
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																			
								15.2	14.7	18.1	17.7	13.6	12.9	13.9	Decreased, 2007-2019	Increased, 2007-2011 Decreased, 2011-2019	No change		
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																			
													5.7	5.2	No linear change	Not available [§]	No change		
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																			
													16.2	13.9	11.3	15.5	No linear change	Not available	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)																	
			31.2			23.4	23.0	21.8	18.6	15.7	11.5	11.5	10.3	9.5	Decreased, 1997-2019	No quadratic change	No change
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											7.8	5.4	4.3	4.7	No linear change	Not available [§]	No change
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											40.4	36.1	37.7	34.8	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Male
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			46.0			30.0	36.7	39.4	33.8	36.4	33.5	34.0	30.8	23.6	Decreased, 1997-2019	No quadratic change	Decreased
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			26.5			11.5	10.3	12.7	10.3	11.6	9.7	8.8	7.8	4.0	Decreased, 1997-2019	No quadratic change	Decreased
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)																	
													9.4	8.5	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male
Injury and Violence

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)																	
			9.6			7.7	11.1	9.6	10.4	8.7	6.6	7.6	9.1	7.9	Decreased, 1997-2019	No quadratic change	No change
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)																	
			41.8			32.4	35.5	32.4	35.6	35.7	28.8	23.6	27.3	24.1	Decreased, 1997-2019	No quadratic change	No change
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)																	
			18.6			12.0	15.9	13.7	13.8	15.1	7.7	10.3	10.3	6.1	Decreased, 1997-2019	No quadratic change	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Male
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †			
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)																			
													5.7	5.0	No linear change	Not available [§]	No change		
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																			
													6.4	6.1	2.2	3.2	Decreased, 2013-2019	Not available	No change
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																			
													7.6	5.3	6.7	5.4	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)																	
								19.7	17.1	18.6	15.5	16.4	19.6		No linear change	No quadratic change	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)																	
								13.1	9.9	9.2	11.8	13.0		No linear change	Not available [§]	No change	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
					23.6	21.8	23.1	21.7	22.1	20.0	20.5	17.6	27.8		No linear change	No quadratic change	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Injury and Violence**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
			18.4			14.3	12.6	12.2	12.4	11.6	10.9	10.6	9.6	14.6	Decreased, 1997-2019	Decreased, 1997-2015 Increased, 2015-2019	Increased
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
			12.7			13.8	9.9	9.9	10.9	12.8	10.9	9.8	9.6	14.0	No linear change	No quadratic change	Increased
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)																	
			6.5			9.8	7.0	5.8	8.4	10.4	5.9	5.7	5.8	6.2	No linear change	No quadratic change	No change
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)																	
			2.5			3.4	2.2	1.4	3.8	4.9	2.6	2.5	2.0	2.9	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Male
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
		78.1				69.6	60.2	62.0	61.1	61.2	49.2	41.4	41.2	31.7	Decreased, 1997-2019	Decreased, 1997-2011 Decreased, 2011-2019	Decreased
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													16.5	14.1	No linear change	Not available [§]	No change
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
		48.4				31.8	26.4	26.2	29.1	26.7	20.3	15.7	14.3	10.3	Decreased, 1997-2019	No quadratic change*	No change
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
		30.0				18.0	15.3	12.5	12.1	12.3	8.4	4.5	4.2	3.4	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)																	
			24.9			14.6	12.3	10.3	9.7	9.6	6.6	3.6	3.3	2.3	Decreased, 1997-2019	No quadratic change	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])																	
												43.9	46.9	52.7	Increased, 2015-2019	Not available [§]	No change
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)																	
												23.4	16.4	27.6	No linear change	Not available	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Tobacco Use**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019									
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)														4.4	4.1	14.2	Increased, 2015-2019			Not available [§]			Increased
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)														3.2	3.0	11.6	Increased, 2015-2019			Not available			Increased
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)														29.3	23.1	30.0	No linear change			Not available			Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Tobacco Use**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019											
QN36: Percentage of students who usually got their own electronic vapor products by buying them in a store (such as a convenience store, supermarket, discount store, gas station, or vape store, during the 30 days before the survey, among students who currently used electronic vapor products and who were aged <18 years)														16.1	17.6	No linear change	Not available [§]	No change							
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)														17.2	10.2	Decreased, 2017-2019	Not available	Decreased							
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)														24.2	21.5	19.6	24.6	23.4	18.3	17.1	13.4	10.5	Decreased, 2003-2019	No change, 2003-2011 Decreased, 2011-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)																	
						40.4	33.2	33.1	36.8	35.1	27.6	24.1	19.0	15.2	Decreased, 2003-2019	No change, 2003-2011 Decreased, 2011-2019	No change
QN39: Percentage of students who tried to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)																	
												43.9	49.9		No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male Alcohol and Other Drug Use															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																	
			37.5			31.9	32.9	28.4	24.4	24.4	21.0	17.9	18.5	17.5	Decreased, 1997-2019	No quadratic change	No change
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)																	
			53.8			46.3	38.0	41.0	40.4	35.6	32.6	25.6	24.4	23.6	Decreased, 1997-2019	No quadratic change	No change
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																	
													12.7	11.8	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male Alcohol and Other Drug Use															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN45: Percentage of students who ever used marijuana (one or more times during their life)																	
			51.9			45.4	36.0	36.0	34.4	39.7	36.9	33.7	30.9	31.9	Decreased, 1997-2019	No quadratic change	No change
QN46: Percentage of students who tried marijuana for the first time before age 13 years																	
			11.3			14.7	13.4	13.4	11.4	13.1	10.6	6.1	8.8	7.7	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)																	
			33.5			22.5	18.1	17.4	19.6	20.6	20.0	17.5	14.6	16.9	Decreased, 1997-2019	Decreased, 1997-2004 Decreased, 2004-2019	No change
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)																	
												8.5	7.6	7.2	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male
Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)																	
													9.5	10.8	No linear change	Not available [§]	No change
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)																	
			9.8			9.8	9.7	9.8	7.1	9.3	6.5	4.5	5.7	4.2	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)																	
		26.5				13.8	14.0	14.7	13.6	12.1	6.7	5.7	7.4	5.1	Decreased, 1997-2019	No quadratic change	No change
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)																	
												3.6	2.2	2.0	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male Alcohol and Other Drug Use																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
						7.2	7.3	8.2	8.2	8.5	5.8	5.8	4.8	4.3	Decreased, 2003-2019	No change, 2003-2011 Decreased, 2011-2019	No change
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
			7.2			7.3	7.3	7.8	5.8	7.6	3.8	5.5	4.2	2.5	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													2.7	2.2	No linear change	Not available [§]	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
			39.8			31.7	21.6	28.8	27.9	26.6	24.8	20.5	23.0	24.8	Decreased, 1997-2019	Decreased, 1997-2004 No change, 2004-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN58: Percentage of students who ever had sexual intercourse																	
		56.9				49.7	48.0	49.0	49.0	51.7	45.9	42.3	40.3	38.3	Decreased, 1997-2019	No quadratic change	No change
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years																	
		10.7				7.6	11.5	10.0	9.3	8.9	7.5	5.4	5.3	4.2	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change
QN60: Percentage of students who had sexual intercourse with four or more persons during their life																	
		23.1				15.8	16.6	14.8	14.7	17.0	14.4	10.6	11.7	9.7	Decreased, 1997-2019	No quadratic change	No change
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)																	
		40.2				34.8	32.5	33.2	31.2	34.6	31.0	31.0	28.8	26.9	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male Sexual Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)																	
			34.8			28.0	24.6	21.4	28.6	22.2	23.8	17.3	17.9	18.3	Decreased, 1997-2019	No quadratic change	No change
QN63: Percentage of students who used a condom during last sexual intercourse (among students who were currently sexually active)																	
			65.5			68.3	69.4	67.5	66.3	55.9	62.1	64.6	53.3	60.5	Decreased, 1997-2019	No quadratic change	No change
QN64: Percentage of students who used birth control pills before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
			15.1			16.6	14.5	14.9	19.4	13.8	15.3	18.7	19.8	22.0	No linear change	No quadratic change	No change
QNIUDIMP: Percentage of students who used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
											2.5	2.6	5.1	4.5	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Sexual Behaviors**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]								
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QNSHPARG: Percentage of students who used a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active))														2.9	1.7	3.0	4.4	No linear change				Not available [§]		No change
QNOTHHPL: Percentage of students who used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)														20.6	23.0	27.9	30.9	Increased, 2013-2019				Not available		No change
QNDUALBC: Percentage of students who used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)														4.6	9.1	7.7	7.9	No linear change				Not available		No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)																	
			10.6			9.2	9.8	11.0	11.1	14.4	11.2	11.6	18.2	11.8	No linear change	No quadratic change	No change
QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)																	
												10.1	12.0	11.4	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						14.3	17.2	17.2	14.8	14.3	13.8	14.9	14.9	15.5	No linear change	No quadratic change	No change
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						19.4	20.4	19.7	20.4	20.6	24.2	20.6	23.0	21.9	No linear change	No quadratic change	No change
QN67: Percentage of students who described themselves as slightly or very overweight																	
		26.3				27.9	29.5	25.0	24.2	23.2	29.3	25.5	28.1	30.4	No linear change	No quadratic change	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)																	
						24.5	27.9	29.7	27.6	27.5	30.0	27.8	30.3	29.9	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)																	
						20.9	21.8	21.2	20.2	23.1	16.5	18.0	20.2	20.0	No linear change	No quadratic change	No change
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						12.1	13.4	12.7	11.4	12.4	10.1	9.3	11.3	11.6	No linear change	No quadratic change	No change
QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						46.9	53.1	47.6	51.8	51.3	55.3	52.7	51.6	48.2	No linear change	Increased, 2003-2015 No change, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						19.3	26.1	22.6	24.1	24.4	27.9	20.1	23.8	20.4	No linear change	Increased, 2003-2013 Decreased, 2013-2019	No change
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)																	
						47.1	49.9	50.8	50.3	50.4	52.0	55.9	54.9	54.3	Increased, 2003-2019	No quadratic change	No change
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)																	
						24.9	25.8	26.1	23.4	26.7	25.3	29.5	37.2	35.8	Increased, 2003-2019	No change, 2003-2009 Increased, 2009-2019	No change
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)																	
						59.3	59.0	59.8	58.7	56.6	56.1	60.7	61.6	62.1	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Male

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
						20.1	23.2	23.5	23.4	26.3	24.4	25.5	27.0	25.5	Increased, 2003-2019	No quadratic change	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						7.5	10.0	8.6	7.9	8.7	7.6	10.4	12.0	12.0	Increased, 2003-2019	No quadratic change	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						53.2	57.6	55.9	57.4	58.4	57.9	55.3	49.2	49.0	Decreased, 2003-2019	No change, 2003-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						22.2	28.0	23.0	22.1	26.3	24.4	21.9	17.2	18.1	Decreased, 2003-2019	No change, 2003-2013 Decreased, 2013-2019	No change
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						11.6	15.2	12.3	10.7	12.8	13.7	10.9	9.0	8.5	Decreased, 2003-2019	No change, 2003-2013 Decreased, 2013-2019	No change
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								15.7	16.3	17.5	20.4	22.7	21.7	23.8	Increased, 2007-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							44.1	39.7	37.1	36.2	33.0	31.9	27.3		Decreased, 2007-2019	No quadratic change	No change
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							34.7	30.1	28.0	26.9	23.9	23.4	19.3		Decreased, 2007-2019	No quadratic change	No change
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)																	
										18.6	18.5	20.0	23.5		Increased, 2013-2019	Not available [§]	No change
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
										39.8	37.9	34.7	30.0		Decreased, 2013-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNMILK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
											14.1	13.2	9.7	10.0	Decreased, 2013-2019	Not available [§]	No change
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)																	
								16.1	17.6	12.6	13.0	14.8	16.3		No linear change	Decreased, 2009-2015 Increased, 2015-2019	No change
QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)																	
								35.0	31.2	42.9	37.3	34.1	29.7		No linear change	Increased, 2009-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Physical Activity**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019												
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															49.5	48.7	44.1	50.5	44.4	No linear change			Not available [§]			Decreased
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															17.2	16.1	13.9	15.5	15.5	No linear change			Not available			No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															30.4	29.5	27.3	31.0	24.8	No linear change			Not available			Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Male
Physical Activity**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																	
						31.3	37.9	27.7	28.6	33.0	24.5	25.3	20.1	18.8	Decreased, 2003-2019	No quadratic change	No change
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																	
								24.6	26.7	36.8	35.4	38.5	41.2	49.5	Increased, 2007-2019	No quadratic change	Increased
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																	
			36.9			43.8	30.2	37.3	41.8	43.9	41.9	44.2	38.2	37.8	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male
Physical Activity

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)																	
			27.4			28.7	20.2	23.5	29.8	24.8	22.5	25.2	25.2	20.9	No linear change	No quadratic change	No change
QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)																	
						55.5	55.2	51.7	51.9	47.9	54.3	53.3	50.5	49.7	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male Other	Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)																	
							68.0	69.6	66.3	66.5	68.2	70.3	70.2			No linear change	No quadratic change	No change
	QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)																	
							2.8	2.3	3.3	2.3	2.1	2.4	1.7			No linear change	No quadratic change	No change
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma																	
							22.6	26.6	24.8	28.9	24.9	25.3	24.9	24.7		No linear change	No quadratic change	No change
	QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)																	
												26.8	23.3	20.7	Decreased, 2015-2019	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Male Other	Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)																	
										60.9	65.7	69.3	71.4	74.8	Increased, 2011-2019	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male
Site-Added

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														57.2	62.3	No linear change	Not available [§]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														46.6	54.5	Increased, 2017-2019	Not available	Increased
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														36.0	40.7	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male
Site-Added

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]										
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019													
QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)															13.3	11.5	12.3	No linear change	Not available [§]	No change							
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)															28.0	14.2	11.2	9.7	10.9	10.9	6.2	6.5	5.2	5.2	Decreased, 1997-2019	Decreased, 1997-2004 Decreased, 2004-2019	No change
QN93: Percentage of students who had oral sex															50.5	52.9	44.5	42.0	40.9	40.0	Decreased, 2009-2019	No quadratic change	No change				
QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)															12.1	10.4	No linear change	Not available	No change								

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Male
Site-Added

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																		
								70.5	69.3	66.8	69.8	64.8	64.4	63.3	Decreased, 2007-2019	No quadratic change	No change	
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																		
								12.4	12.0	14.4	15.1	12.2	11.4	12.1	No linear change	No quadratic change	No change	
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																		
													6.3	6.1	No linear change	Not available [§]	No change	
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																		
												16.7	11.1	11.9	13.6	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Injury and Violence**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 †
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)																	
			15.5			14.1	13.0	13.2	7.8	8.8	8.5	6.6	6.5	5.7	Decreased, 1997-2019	No quadratic change	No change
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											3.4	6.1	3.0	2.9	No linear change	Not available [§]	No change
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											31.6	37.1	30.2	28.6	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Injury and Violence

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			6.6			6.4	9.0	9.0	9.3	8.9	7.6	11.6	9.2	7.2	No linear change	No quadratic change	No change
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			3.7			3.0	3.0	3.0	2.5	3.1	2.7	4.1	1.5	0.9	Decreased, 1997-2019	No quadratic change	No change
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)																	
													4.2	1.4	Decreased, 2017-2019	Not available [§]	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Injury and Violence

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)																	
			3.8			2.3	4.7	6.6	5.2	5.1	3.8	6.7	4.7	5.8	No linear change	No quadratic change	No change
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)																	
			22.8			20.0	23.5	21.5	21.7	21.2	13.4	16.1	14.7	14.5	Decreased, 1997-2019	No quadratic change	No change
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)																	
			5.9			7.8	9.3	7.3	5.1	7.2	4.1	4.8	4.7	5.1	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Injury and Violence

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)																	
												14.2	12.0		No linear change	Not available [§]	No change
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											13.1	13.8	10.2	9.4	Decreased, 2013-2019	Not available	No change
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											11.8	12.1	9.7	8.7	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Injury and Violence**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)																	
									21.9	20.8	24.1	28.6	25.8	27.9	Increased, 2009-2019	No quadratic change	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)																	
									21.9	16.4	25.0	25.1	22.7	No linear change	Not available [§]	No change	
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
					36.7	35.0	35.9	32.1	32.1	31.7	42.5	40.9	46.7	Increased, 2003-2019	No change, 2003-2011 Increased, 2011-2019	Increased	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
			26.1			21.0	18.3	17.8	16.9	18.0	19.2	20.6	19.4	21.3	Decreased, 1997-2019	Decreased, 1997-2007 Increased, 2007-2019	No change
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
			21.4			14.8	13.4	13.8	14.2	14.5	14.3	18.1	15.9	17.2	No linear change	Decreased, 1997-2004 Increased, 2004-2019	No change
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)																	
			9.1			10.3	11.2	9.2	9.1	10.8	9.2	12.9	9.3	9.3	No linear change	No quadratic change	No change
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)																	
			2.2			3.6	3.1	3.2	3.2	4.1	3.2	5.3	3.4	2.6	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
		76.3				72.4	58.7	62.4	57.1	57.0	44.9	46.8	39.8	29.3	Decreased, 1997-2019	Decreased, 1997-2011 Decreased, 2011-2019	Decreased
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													14.2	9.0	Decreased, 2017-2019	Not available [§]	Decreased
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
		45.3				33.4	26.0	25.8	23.1	21.4	15.5	18.0	14.0	7.2	Decreased, 1997-2019	No quadratic change*	Decreased
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
		24.8				18.8	13.5	14.2	12.0	10.5	6.3	6.8	5.4	2.4	Decreased, 1997-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)																	
			19.6			15.2	9.9	10.8	8.9	8.1	4.6	5.5	3.9	1.9	Decreased, 1997-2019	No quadratic change	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])																	
												39.5	41.7	54.5	Increased, 2015-2019	Not available [§]	Increased
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)																	
												23.3	11.3	24.0	No linear change	Not available	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)																	
												2.0	1.2	7.5	Increased, 2015-2019	Not available [§]	Increased
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)																	
												1.1	0.8	5.2	Increased, 2015-2019	Not available	Increased
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)																	
												29.4	18.4	25.7	No linear change	Not available	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Tobacco Use**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019											
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)														3.1	2.0	No linear change	Not available [§]	No change							
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)														12.6	9.0	11.1	9.5	11.0	7.6	10.2	7.6	4.6	Decreased, 2003-2019	No quadratic change	Decreased
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)														37.4	28.3	29.1	26.1	24.8	19.5	22.2	16.4	10.1	Decreased, 2003-2019	No quadratic change	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN39: Percentage of students who tried to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)																	
												48.5	52.3	No linear change	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																	
			22.6			21.3	24.7	21.3	18.9	16.1	16.8	13.5	15.3	13.0	Decreased, 1997-2019	No quadratic change	No change
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)																	
			44.5			44.2	36.8	40.1	35.2	33.3	28.0	31.2	28.6	23.1	Decreased, 1997-2019	No quadratic change	Decreased
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																	
													13.8	8.6	Decreased, 2017-2019	Not available [§]	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female Alcohol and Other Drug Use															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN45: Percentage of students who ever used marijuana (one or more times during their life)																	
			44.0			41.1	32.6	33.9	28.5	34.8	30.6	32.6	32.9	31.6	Decreased, 1997-2019	Decreased, 1997-2009 No change, 2009-2019	No change
QN46: Percentage of students who tried marijuana for the first time before age 13 years																	
			6.8			8.2	6.4	6.9	6.9	6.6	6.0	6.3	5.8	6.1	No linear change	No quadratic change	No change
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)																	
			23.3			19.5	13.4	15.4	12.5	17.4	15.3	16.9	16.7	14.8	Decreased, 1997-2019	Decreased, 1997-2004 No change, 2004-2019	No change
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)																	
												11.6	7.5	4.4	Decreased, 2015-2019	Not available [§]	Decreased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female																			
Alcohol and Other Drug Use																			
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)																			
															11.9	10.9	No linear change	Not available [§]	No change
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)																			
			6.3			9.3	6.9	7.0	5.0	5.0	2.0	4.2	3.1	2.0	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change		
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)																			
			22.5			14.6	13.0	13.4	13.2	10.3	7.2	8.6	5.0	5.4	Decreased, 1997-2019	No quadratic change	No change		
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)																			
												3.1	1.6	0.8	Decreased, 2015-2019	Not available	No change		

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
						5.9	4.4	4.4	5.5	4.1	2.1	4.1	2.8	2.0	Decreased, 2003-2019	No quadratic change	No change
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
			4.6			6.7	4.0	4.1	2.2	2.4	1.5	3.0	2.2	1.1	Decreased, 1997-2019	No quadratic change	No change
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													2.0	1.4	No linear change	Not available [§]	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
			28.2			28.9	17.9	25.1	23.1	22.1	15.9	21.0	21.3	22.6	Decreased, 1997-2019	Decreased, 1997-2013 Increased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN58: Percentage of students who ever had sexual intercourse																	
			50.3			54.3	44.6	51.5	47.8	51.9	43.3	41.4	36.5	40.4	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years																	
			3.4			3.7	4.1	5.8	4.0	5.5	3.2	4.1	1.8	2.1	Decreased, 1997-2019	Increased, 1997-2007 Decreased, 2007-2019	No change
QN60: Percentage of students who had sexual intercourse with four or more persons during their life																	
			12.6			13.7	10.6	13.9	10.7	16.3	10.5	10.2	7.6	7.2	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)																	
			38.3			42.4	34.5	39.6	36.1	40.9	32.3	29.6	28.9	30.7	Decreased, 1997-2019	No change, 1997-2011 Decreased, 2011-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female Sexual Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)																	
			19.4			17.1	14.4	17.2	18.0	12.4	13.6	16.4	17.3	11.9	No linear change	No quadratic change	No change
QN63: Percentage of students who used a condom during last sexual intercourse (among students who were currently sexually active)																	
			49.7			56.9	61.4	52.6	54.5	46.4	45.1	43.8	44.3	45.6	Decreased, 1997-2019	Increased, 1997-2004 Decreased, 2004-2019	No change
QN64: Percentage of students who used birth control pills before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
			22.3			18.3	22.2	25.0	26.9	27.1	24.2	26.1	27.9	26.5	Increased, 1997-2019	No quadratic change	No change
QNIUDIMP: Percentage of students who used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
											2.7	7.8	10.1	8.5	Increased, 2013-2019	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Sexual Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019†
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSHPARG: Percentage of students who used a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active))															No linear change	Not available§	No change
QNOTHHPL: Percentage of students who used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)															Increased, 2013-2019	Not available	No change
QNDUALBC: Percentage of students who used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)															Increased, 2013-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)																	
			17.2			17.4	12.5	15.8	14.8	17.7	18.6	17.5	15.4	15.2	No linear change	No quadratic change	No change
QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)																	
												12.9	10.3	10.1	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						16.1	16.5	15.2	16.0	16.6	17.0	19.2	17.1	20.3	Increased, 2003-2019	No quadratic change	No change
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]																	
						9.3	10.2	10.8	14.1	12.1	11.3	16.2	17.3	14.6	Increased, 2003-2019	No quadratic change	No change
QN67: Percentage of students who described themselves as slightly or very overweight																	
		38.0				39.9	37.2	37.5	36.5	36.7	35.5	38.8	39.6	39.3	No linear change	No quadratic change	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)																	
						27.0	26.5	27.9	24.7	25.6	27.2	28.6	32.3	35.6	Increased, 2003-2019	No change, 2003-2011 Increased, 2011-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

†Based on t-test analysis, $p < 0.05$.

§Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)																	
						16.2	19.8	16.6	17.4	15.1	11.5	11.3	12.9	16.4	Decreased, 2003-2019	No quadratic change	No change
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						7.8	9.2	8.5	8.0	7.0	5.8	5.8	6.2	9.0	No linear change	No quadratic change	Increased
QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						44.6	46.8	46.3	49.9	49.1	55.7	52.7	45.1	46.7	No linear change	Increased, 2003-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female																	
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019[†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)															No linear change	Increased, 2003-2009 Decreased, 2009-2019	No change
						19.0	21.3	19.9	24.5	21.5	23.5	21.9	17.5	21.6			
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)															Increased, 2003-2019	No quadratic change	No change
						33.3	44.3	42.0	40.7	40.8	41.9	43.2	45.1	46.8			
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)															Increased, 2003-2019	No change, 2003-2009 Increased, 2009-2019	Increased
						26.3	22.6	22.5	22.7	24.8	27.8	28.2	33.7	39.5			
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)															No linear change	Decreased, 2003-2015 Increased, 2015-2019	No change
						59.5	60.6	60.5	58.7	61.5	56.0	55.7	62.2	64.0			

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female															Linear Change*	Quadratic Change*	Change from 2017-2019 †
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
						15.8	15.1	17.7	19.8	19.4	17.6	18.0	20.5	22.9	Increased, 2003-2019	No quadratic change	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						5.3	4.8	4.8	4.6	5.2	4.8	6.5	8.5	9.9	Increased, 2003-2019	No change, 2003-2013 Increased, 2013-2019	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						57.9	58.5	58.9	60.0	55.2	56.5	58.8	51.9	47.4	Decreased, 2003-2019	No change, 2003-2015 Decreased, 2015-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female															Linear Change*	Quadratic Change*	Change from 2017-2019 †
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						21.9	24.3	20.9	22.6	24.5	20.7	24.0	18.9	18.9	Decreased, 2003-2019	No quadratic change	No change
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						8.8	11.7	9.9	11.2	11.9	9.3	11.1	8.4	8.6	No linear change	No quadratic change	No change
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								17.2	20.0	17.8	23.2	24.7	24.7	27.2	Increased, 2007-2019	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Female															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
Weight Management and Dietary Behaviors																	
Health Risk Behavior and Percentages																	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								36.9	31.6	35.7	29.6	31.6	24.3	23.5	Decreased, 2007-2019	No quadratic change	No change
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								28.2	23.3	24.7	22.0	22.5	17.5	15.5	Decreased, 2007-2019	No quadratic change	No change
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)																	
										25.5	26.5	32.0	37.0	Increased, 2013-2019	Not available [§]	Increased	
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
								28.8	22.3	21.3	18.7	Decreased, 2013-2019	Not available	No change			

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female															Linear Change*	Quadratic Change*	Change from 2017-2019 †		
Weight Management and Dietary Behaviors																			
Health Risk Behavior and Percentages																			
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QNMILK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																			
											6.4	3.9	4.5	4.3	No linear change	Not available [§]	No change		
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)																			
											13.5	13.4	11.9	12.4	16.1	16.4	No linear change	No quadratic change	No change
QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)																			
											28.8	28.6	37.2	32.2	26.1	23.4	Decreased, 2009-2019	Increased, 2009-2013 Decreased, 2013-2019	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Physical Activity**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															29.0	30.8	29.9	30.8	30.3	No linear change			Not available [§]	No change
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															23.9	23.6	18.8	22.9	22.5	No linear change			Not available	No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															13.3	15.4	12.9	12.8	13.0	No linear change			Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Physical Activity**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																	
						30.5	32.9	27.1	29.2	31.0	28.8	25.4	22.2	24.1	Decreased, 2003-2019	No quadratic change	No change
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																	
								18.1	19.2	25.6	33.6	41.9	41.6	45.6	Increased, 2007-2019	Increased, 2007-2015 No change, 2015-2019	No change
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																	
			25.5			25.8	20.1	24.4	23.7	26.5	26.7	31.1	23.7	24.5	No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

Female
Physical Activity

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)																	
			16.9			19.0	14.3	16.6	16.1	15.3	16.1	19.3	13.1	14.7	No linear change	No quadratic change	No change
QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)																	
					46.4	49.7	45.3	44.6	44.6	47.3	48.2	46.5	42.0		No linear change	No quadratic change	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Other

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)																		
								71.6	68.4	69.8	72.5	71.5	76.8	73.4	Increased, 2007-2019	No quadratic change	No change	
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)																		
								2.6	1.8	2.7	2.2	2.1	1.8	2.1	No linear change	No quadratic change	No change	
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma																		
								21.6	25.4	23.8	24.4	21.1	25.6	26.9	22.4	No linear change	No quadratic change	Decreased
QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)																		
												21.8	20.8	18.3	No linear change	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female Other	Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)																	
										74.2	80.8	78.6	83.2	82.3	Increased, 2011-2019	Not available [§]	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Female
Site-Added

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												62.2	63.6		No linear change	Not available [§]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												53.0	54.7		No linear change	Not available	No change
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												38.8	41.0		No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Site-Added**

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]										
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019													
QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)															28.0	22.5	22.6	No linear change	Not available [§]	No change							
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)															22.1	14.2	8.2	9.4	7.9	7.5	3.8	7.2	5.0	3.2	Decreased, 1997-2019	No quadratic change	No change
QN93: Percentage of students who had oral sex															47.8	49.6	38.8	41.5	37.2	38.7				Decreased, 2009-2019	No quadratic change	No change	
QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)															5.5	7.1				No linear change	Not available	No change					

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Female
Site-Added**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]			
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																			
								83.6	85.8	82.6	85.5	81.7	78.1	78.6	Decreased, 2007-2019	No quadratic change	No change		
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																			
								18.1	17.7	22.1	20.7	15.0	14.8	15.8	Decreased, 2007-2019	No change, 2007-2011 Decreased, 2011-2019	No change		
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																			
													4.4	3.7	No linear change	Not available [§]	No change		
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																			
													15.3	16.5	10.6	17.1	No linear change	Not available	Increased

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)																	
			22.4			17.5	17.4	16.5	12.8	11.0	8.6	7.6	7.1	6.5	Decreased, 1997-2019	No quadratic change	No change
QN10: Percentage of students who drove a car or other vehicle when they had been drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											5.7	4.8	3.8	4.0	No linear change	Not available [¶]	No change
QN11: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)																	
											36.0	37.3	34.8	31.3	Decreased, 2013-2019	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			26.4			18.9	23.5	25.4	22.5	21.9	22.3	23.5	20.9	16.1	Decreased, 1997-2019	No quadratic change	No change
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
			15.5			7.4	6.7	8.2	6.2	6.8	6.3	5.8	4.4	2.6	Decreased, 1997-2019	No quadratic change	No change
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)																	
													5.4	4.2	No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)																	
			6.5			4.6	7.7	7.4	7.3	5.6	5.1	5.1	6.3	6.3	No linear change	No quadratic change	No change
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)																	
			32.1			25.9	28.3	25.7	27.7	26.5	19.1	17.8	19.8	17.4	Decreased, 1997-2019	No quadratic change	No change
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)																	
			12.1			9.6	11.7	9.5	8.8	10.6	5.5	6.6	6.3	4.5	Decreased, 1997-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [‡]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)																	
												8.6	8.4		No linear change	Not available [¶]	No change
QN21: Percentage of students who experienced sexual dating violence (being forced by someone they were dating or going out with to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											8.6	8.8	5.9	6.7	Decreased, 2013-2019	Not available	No change
QN22: Percentage of students who experienced physical dating violence (being physically hurt on purpose by someone they were dating or going out with [counting such things as being hit, slammed into something, or injured with an object or weapon] one or more times during the 12 months before the survey, among students who dated or went out with someone during the 12 months before the survey)																	
											8.4	8.0	7.9	7.5	No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)																	
									21.3	19.0	21.7	23.4	21.6	25.0	Increased, 2009-2019	No quadratic change	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)																	
										17.5	13.3	17.8	19.3	18.8	Increased, 2011-2019	Not available [¶]	No change
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
					30.1	28.5	29.0	26.6	27.0	24.5	30.6	29.1	36.7	Increased, 2003-2019	Decreased, 2003-2013 Increased, 2013-2019	Increased	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Injury and Violence

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
			22.3			17.9	15.7	14.7	14.1	14.2	14.6	14.9	14.4	18.1	Decreased, 1997-2019	Decreased, 1997-2009 No change, 2009-2019	No change
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
			17.1			14.3	11.6	11.6	11.7	13.8	11.8	13.1	12.5	15.3	No linear change	Decreased, 1997-2007 No change, 2007-2019	No change
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)																	
			7.6			9.3	8.9	7.1	8.2	9.2	6.7	8.1	6.3	7.0	No linear change	No quadratic change	No change
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)																	
			2.3			2.9	2.7	2.2	3.3	3.8	2.3	3.5	2.0	2.3	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Tobacco Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [‡]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
		77.0				71.3	60.9	62.2	59.1	59.7	46.8	44.5	41.5	31.6	Decreased, 1997-2019	Decreased, 1997-2011 Decreased, 2011-2019	Decreased
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													15.0	11.4	Decreased, 2017-2019	Not available [¶]	Decreased
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
		47.7				34.2	27.4	27.0	27.1	25.1	18.9	17.0	15.1	9.7	Decreased, 1997-2019	No quadratic change	Decreased
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
		28.6				19.7	15.0	14.3	12.8	11.9	7.8	5.5	5.5	3.2	Decreased, 1997-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Tobacco Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)																	
		23.5				16.0	11.8	11.3	10.1	9.0	5.9	4.7	4.2	2.3	Decreased, 1997-2019	No quadratic change	No change
QN33: Percentage of students who smoked more than 10 cigarettes per day (on the days they smoked during the 30 days before the survey, among students who currently smoked cigarettes)																	
		26.7				17.0	18.9	14.2	14.8	16.8	11.0	6.5	6.8	9.2	Decreased, 1997-2019	No quadratic change	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])																	
												40.6	45.0	55.7	Increased, 2015-2019	Not available [¶]	Increased

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**White*
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)															22.8	14.4	28.1	No linear change			Not available [¶]			Increased
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)															3.4	2.8	12.8	Increased, 2015-2019			Not available			Increased
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)															2.1	2.0	9.8	Increased, 2015-2019			Not available			Increased
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)															28.8	21.8	30.0	No linear change			Not available			Increased

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Tobacco Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [‡]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN36: Percentage of students who usually got their own electronic vapor products by buying them in a store (such as a convenience store, supermarket, discount store, gas station, or vape store, during the 30 days before the survey, among students who currently used electronic vapor products and who were aged <18 years)																	
													12.1	12.2	No linear change	Not available [¶]	No change
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)																	
													10.9	6.8	Decreased, 2017-2019	Not available	Decreased
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)																	
						18.4	14.9	14.7	17.0	16.4	11.9	12.2	10.9	7.3	Decreased, 2003-2019	No change, 2003-2011 Decreased, 2011-2019	Decreased

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Tobacco Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)																	
					39.9	31.3	31.3	32.3	30.2	23.2	22.2	18.5	13.1		Decreased, 2003-2019	Decreased, 2003-2011 Decreased, 2011-2019	Decreased
QN39: Percentage of students who tried to quit using all tobacco products (including cigarettes, cigars, smokeless tobacco, shisha or hookah tobacco, and electronic vapor products, ever during the 12 months before the survey, among students who used any tobacco products during the 12 months before the survey)																	
												48.0	51.4		No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																	
		29.3				26.4	28.8	24.8	21.5	19.4	17.5	14.3	16.1	14.7	Decreased, 1997-2019	No quadratic change	No change
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)																	
		49.1				46.7	38.6	41.0	38.5	35.2	30.5	29.1	28.7	23.9	Decreased, 1997-2019	No quadratic change	Decreased
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																	
													14.4	11.4	Decreased, 2017-2019	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN45: Percentage of students who ever used marijuana (one or more times during their life)																	
			46.2			43.6	33.7	33.9	30.4	35.8	31.7	31.8	29.9	29.5	Decreased, 1997-2019	Decreased, 1997-2007 Decreased, 2007-2019	No change
QN46: Percentage of students who tried marijuana for the first time before age 13 years																	
			8.3			10.9	9.6	9.3	8.7	8.4	7.0	5.3	6.4	5.8	Decreased, 1997-2019	No change, 1997-2004 Decreased, 2004-2019	No change
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)																	
			26.8			21.4	15.4	15.3	15.2	17.8	16.0	16.1	14.6	14.5	Decreased, 1997-2019	Decreased, 1997-2007 No change, 2007-2019	No change
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)																	
												9.1	7.2	5.3	Decreased, 2015-2019	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [‡]	Change from 2017-2019 [§]													
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019															
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)														9.7	9.9	No linear change	Not available [¶]	No change											
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)														8.4	9.4	8.4	8.3	5.9	6.6	4.2	3.1	3.5	2.8	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change			
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)														25.9	14.4	14.0	14.2	13.3	10.4	6.6	6.3	5.6	5.2	Decreased, 1997-2019	No quadratic change	No change			
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)														2.3	0.9	1.1											Decreased, 2015-2019	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[‡]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
						6.5	5.9	5.8	6.5	5.5	3.5	4.1	3.2	2.6	Decreased, 2003-2019	No quadratic change	No change
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
			5.8			6.8	5.8	5.5	3.6	4.2	2.2	2.9	2.5	1.5	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													1.7	1.3	No linear change	Not available [¶]	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
			34.2			31.0	19.7	26.9	25.1	23.4	18.7	20.1	21.0	23.5	Decreased, 1997-2019	Decreased, 1997-2015 No change, 2015-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN58: Percentage of students who ever had sexual intercourse																	
			50.8			51.4	43.9	48.3	46.6	50.4	43.5	40.8	37.5	38.7	Decreased, 1997-2019	No quadratic change	No change
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years																	
			5.6			5.0	6.8	6.5	5.5	5.4	4.2	3.4	2.9	2.5	Decreased, 1997-2019	No change, 1997-2007 Decreased, 2007-2019	No change
QN60: Percentage of students who had sexual intercourse with four or more persons during their life																	
			15.5			13.6	12.0	12.4	11.8	14.7	11.4	8.9	9.0	8.0	Decreased, 1997-2019	No quadratic change	No change
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)																	
			37.0			38.4	32.2	35.2	32.5	37.0	32.0	30.1	28.3	28.6	Decreased, 1997-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White* Sexual Behaviors		Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN62: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)																		
		27.4				21.9	19.9	19.5	22.4	15.2	16.8	14.4	17.5	14.5	Decreased, 1997-2019	No quadratic change	No change	
QN63: Percentage of students who used a condom during last sexual intercourse (among students who were currently sexually active)																		
		53.9				61.1	64.4	59.2	58.7	50.9	53.6	55.0	47.0	52.8	Decreased, 1997-2019	Increased, 1997-2004 Decreased, 2004-2019	No change	
QN64: Percentage of students who used birth control pills before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																		
		19.9				18.7	19.5	21.2	24.1	21.3	21.4	24.0	28.1	25.9	Increased, 1997-2019	No quadratic change	No change	
QNIUDIMP: Percentage of students who used an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																		
											2.7	5.2	8.8	5.8	Increased, 2013-2019	Not available [¶]	No change	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSHPARG: Percentage of students who used a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active))																	
											4.5	5.8	5.7	7.0	No linear change	Not available [¶]	No change
QNOTHHPL: Percentage of students who used birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
											28.6	35.1	42.6	38.6	Increased, 2013-2019	Not available	No change
QNDUALBC: Percentage of students who used both a condom during last sexual intercourse and birth control pills; an IUD (such as Mirena or ParaGard) or implant (such as Implanon or Nexplanon); or a shot (such as Depo-Provera), patch (such as OrthoEvra), or birth control ring (such as NuvaRing) before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)																	
											6.4	12.3	13.0	12.4	Increased, 2013-2019	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNBCNONE: Percentage of students who did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)																	
			14.0			12.7	11.1	12.6	12.6	15.3	13.0	13.9	15.3	13.7	No linear change	No quadratic change	No change
QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)																	
												10.1	10.0	10.2	No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [†]																	
				15.0		16.4	15.4	15.2	15.2	15.7	16.2	15.6	18.7		No linear change	No quadratic change	No change
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [†]																	
					14.2	15.4	15.1	17.2	16.1	17.7	18.1	19.9	18.3		Increased, 2003-2019	No quadratic change	No change
QN67: Percentage of students who described themselves as slightly or very overweight																	
		32.2				34.9	34.4	32.0	30.7	30.6	32.8	33.3	34.3	36.5	No linear change	No quadratic change	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)																	
					25.7	28.1	29.3	27.6	26.8	29.5	28.3	31.5	34.3		Increased, 2003-2019	No change, 2003-2015 Increased, 2015-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)																	
						17.7	20.7	18.6	18.3	18.3	13.5	14.2	16.4	18.4	Decreased, 2003-2019	No quadratic change	No change
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						9.7	11.4	10.3	9.9	9.3	6.9	7.3	8.2	10.6	Decreased, 2003-2019	Decreased, 2003-2015 Increased, 2015-2019	Increased
QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						46.3	49.1	46.2	49.7	49.0	54.2	52.1	48.1	46.4	No linear change	Increased, 2003-2015 Decreased, 2015-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						19.4	22.4	20.4	23.2	22.2	24.5	20.3	19.5	19.0	No linear change	Increased, 2003-2013 Decreased, 2013-2019	No change
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)																	
						38.6	46.7	46.1	45.4	44.8	46.7	48.8	50.3	50.7	Increased, 2003-2019	No quadratic change	No change
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)																	
						24.7	22.8	22.5	20.8	24.6	25.7	27.7	33.3	36.7	Increased, 2003-2019	No change, 2003-2009 Increased, 2009-2019	No change
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)																	
						59.3	58.7	59.7	57.4	58.2	54.6	56.5	61.8	62.5	No linear change	No change, 2003-2013 Increased, 2013-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
						17.4	19.2	20.1	20.7	21.8	21.0	20.3	22.6	24.3	Increased, 2003-2019	No quadratic change	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						5.9	6.9	6.2	6.0	6.1	5.7	7.3	9.3	11.0	Increased, 2003-2019	No change, 2003-2013 Increased, 2013-2019	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						56.1	58.5	58.4	59.7	56.6	57.4	59.0	50.8	48.1	Decreased, 2003-2019	No change, 2003-2015 Decreased, 2015-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						21.9	26.2	21.5	22.5	24.9	22.2	23.7	17.9	18.0	Decreased, 2003-2019	No change, 2003-2015 Decreased, 2015-2019	No change
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						10.4	13.0	10.5	10.4	11.7	10.9	10.8	8.7	7.9	Decreased, 2003-2019	No quadratic change	No change
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
								16.0	17.8	16.8	21.5	23.2	22.4	24.3	Increased, 2007-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							42.2	37.5	38.2	35.0	33.6	30.0	27.8	Decreased, 2007-2019	No quadratic change	No change	
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							32.7	28.0	27.8	26.1	24.1	21.9	18.6	Decreased, 2007-2019	No quadratic change	No change	
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)																	
										20.5	19.7	24.9	30.1	Increased, 2013-2019	Not available [¶]	Increased	
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
										36.5	32.8	29.9	25.3	Decreased, 2013-2019	Not available	Decreased	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNMLK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
											10.9	9.4	7.3	7.1	Decreased, 2013-2019	Not available [¶]	No change
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)																	
								14.6	14.6	11.2	12.0	14.0	16.0	No linear change	Decreased, 2009-2013 Increased, 2013-2019	No change	
QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)																	
								33.0	30.5	42.1	36.1	31.5	27.8	Decreased, 2009-2019	Increased, 2009-2013 Decreased, 2013-2019	No change	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Physical Activity

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															40.1	41.6	37.9	42.6	38.1	No linear change			Not available [¶]	Decreased
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															19.4	18.4	15.3	17.2	18.2	No linear change			Not available	No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															22.5	23.6	20.0	23.4	19.2	No linear change			Not available	Decreased

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

White*
Physical Activity

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																	
						28.5	32.8	25.5	26.7	30.4	25.1	24.4	19.6	20.3	Decreased, 2003-2019	No quadratic change	No change
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																	
								20.3	22.8	30.9	33.5	41.7	42.4	48.3	Increased, 2007-2019	No quadratic change	Increased
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																	
			29.9			34.9	25.0	29.2	32.8	33.9	33.6	36.1	30.0	28.9	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*
Physical Activity

Health Risk Behavior and Percentages

Linear Change[†]

Quadratic Change[†]

Change from
2017-2019[§]

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)

22.0

24.5

17.9

20.3

24.2

20.3

19.3

22.1

19.6

17.5

No linear change

No quadratic change

No change

QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)

50.7

52.5

47.1

47.7

46.1

50.1

49.9

48.0

45.8

No linear change

No quadratic change

No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White* Other	Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)																	
								71.7	71.2	69.4	70.3	73.1	75.8	75.0		Increased, 2007-2019	No quadratic change	No change
	QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)																	
								2.2	1.7	2.5	1.7	1.4	1.9	1.3		No linear change	No quadratic change	No change
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma																	
								21.6	24.7	23.8	25.1	22.0	24.5	26.0	22.3	No linear change	No quadratic change	No change
	QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)																	
												25.7	22.6	20.2	Decreased, 2015-2019	Not available [¶]	No change	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

White*
Other

Health Risk Behavior and Percentages

Linear Change[†]

Quadratic Change[†]

Change from
2017-2019[§]

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)

69.3 74.9 77.6 78.8 81.2 Increased, 2011-2019 Not available[¶] No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**White*
Site-Added**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														61.6	63.5	No linear change	Not available [¶]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														50.8	54.5	No linear change	Not available	No change
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														37.5	40.8	No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey

Trend Analysis Report

White*
Site-Added

Health Risk Behavior and Percentages

Linear Change[†]

Quadratic Change[†]

Change from
2017-2019[§]

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)

19.8 16.9 17.5 No linear change Not available[¶] No change

QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)

26.3 15.0 10.0 10.0 9.8 9.4 5.0 5.1 4.8 4.1 Decreased, 1997-2019 No quadratic change No change

QN93: Percentage of students who had oral sex

49.7 50.9 41.5 42.5 39.4 40.1 Decreased, 2009-2019 No quadratic change No change

QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)

7.7 8.4 No linear change Not available No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**White*
Site-Added**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																	
								77.9	77.9	75.7	78.6	75.1	72.8	71.7	Decreased, 2007-2019	No quadratic change	No change
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																	
								15.4	13.9	17.3	17.8	12.8	12.2	13.0	Decreased, 2007-2019	Increased, 2007-2013 Decreased, 2013-2019	No change
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																	
													3.6	4.1	No linear change	Not available [¶]	No change
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																	
											14.6	13.3	11.5	15.3	No linear change	Not available	Increased

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Injury and Violence

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)																	
							22.9	24.3	17.0	15.4	16.1	13.7	15.5	14.1	Decreased, 2004-2019	No quadratic change	No change
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
							17.8	13.5	10.8	25.5	7.9	13.4	14.5	9.0	No linear change	No quadratic change	No change
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
							6.9	5.1	5.1	8.3	4.3	9.5	5.8	2.8	No linear change	No quadratic change	No change
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)																	
													11.7	7.1	No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Injury and Violence

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)																	
							8.8	9.7	8.9	8.9	3.4	18.7	9.7	10.3	No linear change	No quadratic change	No change
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)																	
							38.0	34.8	33.0	38.9	32.2	31.4	28.3	28.7	Decreased, 2004-2019	No quadratic change	No change
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)																	
							18.5	14.8	9.9	13.9	6.3	15.9	10.6	9.5	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Injury and Violence**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]						
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019								
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)														11.7	9.2	No linear change	Not available [¶]	No change				
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)														14.8	14.5	19.1	12.7	16.8	17.8	No linear change	No quadratic change	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)														14.0	10.7	10.3	11.3	15.4	No linear change	Not available	No change	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Injury and Violence

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
							21.8	30.9	23.0	26.1	30.6	37.5	28.1	36.6	Increased, 2004-2019	No quadratic change	No change
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
							9.9	14.7	14.7	17.8	14.9	17.3	9.7	21.0	No linear change	No quadratic change	Increased
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
							10.2	10.6	13.6	12.9	14.2	17.5	13.4	19.3	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Tobacco Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
							47.4	61.5	55.7	53.9	45.9	35.5	27.8	20.9	Decreased, 2004-2019	No change, 2004-2011 Decreased, 2011-2019	No change
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													15.3	11.7	No linear change	Not available [¶]	No change
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
							14.9	14.4	17.2	14.4	11.9	10.6	6.3	4.9	Decreased, 2004-2019	No quadratic change	No change
QNFRDIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
							8.1	5.2	4.6	5.7	4.6	4.2	0.3	2.3	Decreased, 2004-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Tobacco Use**

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)																	
						4.7	3.3	2.6	4.7	3.3	1.9	0.3	2.3		Decreased, 2004-2019	No quadratic change	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])																	
												42.3	41.7	40.2	No linear change	Not available [¶]	No change
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)																	
												24.7	7.3	13.0	Decreased, 2015-2019	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Tobacco Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)														Not available	Not available [¶]	No change	
												0.0	0.3	3.0			
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)														Not available	Not available	No change	
												0.0	0.3	3.0			
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)														Decreased, 2015-2019	Not available	No change	
												28.8	11.4	15.1			

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Tobacco Use**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]								
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)														6.4	2.3	No linear change	Not available [¶]	No change						
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)														18.2	17.3	16.7	20.0	18.8	17.7	7.8	8.7	Decreased, 2004-2019	No change, 2004-2015 Decreased, 2015-2019	No change
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)														24.9	25.5	23.9	25.6	25.7	23.0	12.3	9.9	Decreased, 2004-2019	No change, 2004-2015 Decreased, 2015-2019	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																	
						28.1	21.1	19.6	20.9	26.6	17.4	16.6	14.6		Decreased, 2004-2019	No quadratic change	No change
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																	
												5.3	4.8		No linear change	Not available [¶]	No change
QN45: Percentage of students who ever used marijuana (one or more times during their life)																	
						40.0	42.1	37.7	43.4	46.0	40.4	44.6	40.8		No linear change	No quadratic change	No change
QN46: Percentage of students who tried marijuana for the first time before age 13 years																	
						13.3	13.7	12.3	14.3	16.5	9.3	11.2	11.3		No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)																	
						18.9	23.9	21.3	22.9	27.5	22.1	20.7	22.2		No linear change	No quadratic change	No change
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)																	
											13.1	6.9	7.0		Decreased, 2015-2019	Not available [¶]	No change
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)																	
												12.5	12.8		No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)																	
						5.7	6.3	3.2	6.3	2.6	8.2	4.0	3.1		No linear change	No quadratic change	No change
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)																	
						6.2	10.2	9.6	12.6	7.5	9.1	8.9	5.9		No linear change	No change, 2004-2011 No change, 2011-2019	No change
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)																	
											9.8	3.7	2.6		Decreased, 2015-2019	Not available [¶]	No change
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
						4.0	8.8	5.7	6.9	5.1	8.4	7.1	5.9		No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*

Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
							3.5	6.9	4.4	5.7	3.7	13.3	5.9	3.1	No linear change	Increased, 2004-2015 Decreased, 2015-2019	No change
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													3.3	4.6	No linear change	Not available [¶]	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
							18.5	22.6	27.5	27.7	27.6	24.0	29.4	24.2	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Sexual Behaviors

Health Risk Behavior and Percentages

Linear Change†

Quadratic Change†

Change from
2017-2019 §

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)

15.4 16.5 14.5 No linear change Not available¶ No change

*Non-Hispanic.

†Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

§Based on t-test analysis, $p < 0.05$.

¶Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [†]																	
							20.7	23.1	16.6	14.8	13.3	22.9	20.0	13.6	No linear change	No quadratic change	No change
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [†]																	
							15.5	17.0	21.9	19.5	19.1	17.7	26.4	22.5	Increased, 2004-2019	No quadratic change	No change
QN67: Percentage of students who described themselves as slightly or very overweight																	
							23.5	25.1	26.1	24.8	31.6	21.0	33.1	25.2	No linear change	No quadratic change	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)																	
							20.0	24.7	14.1	26.5	25.9	24.1	31.2	24.7	Increased, 2004-2019	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages

Linear Change[†]

Quadratic Change[†]

Change from
2017-2019[§]

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)

22.8 22.6 19.7 25.8 18.6 17.8 20.9 23.0 No linear change No quadratic change No change

QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)

11.9 12.8 7.9 11.9 15.1 9.2 14.0 10.8 No linear change No quadratic change No change

QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)

56.9 51.2 59.7 52.9 56.8 61.4 50.1 48.7 No linear change No quadratic change No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
						34.5	29.6	30.3	27.7	29.7	25.5	24.5	23.9	Decreased, 2004-2019	No quadratic change	No change	
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)																	
						51.8	52.8	45.8	53.8	49.2	54.8	48.7	52.1	No linear change	No quadratic change	No change	
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)																	
						35.3	37.0	39.1	33.4	32.7	35.5	50.0	44.9	Increased, 2004-2019	No quadratic change	No change	
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)																	
						71.2	66.9	68.6	68.2	65.5	69.9	68.1	69.2	No linear change	No quadratic change	No change	

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
							20.5	25.4	27.2	28.6	22.8	34.8	36.0	26.9	Increased, 2004-2019	No quadratic change	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
							12.0	10.9	7.7	11.4	10.0	17.6	18.0	11.6	Increased, 2004-2019	No quadratic change	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
							52.3	46.3	51.6	54.6	55.3	45.5	45.7	41.4	No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						24.7	23.5	19.8	25.6	24.0	15.4	16.7	18.4	Decreased, 2004-2019	No quadratic change	No change	
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
						15.8	14.5	12.6	13.9	16.7	10.0	7.8	9.7	Decreased, 2004-2019	No quadratic change	No change	
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
						18.2	19.5	24.2	20.9	22.4	25.6	29.4	Increased, 2007-2019	No quadratic change	No change		

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							29.1	24.9	24.3	25.3	28.6	21.4	19.6		No linear change	No quadratic change	No change
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)																	
							23.8	18.4	16.1	20.2	20.5	15.1	15.9		No linear change	No quadratic change	No change
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)																	
										34.9	41.0	34.5	32.2		No linear change	Not available [¶]	No change
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)																	
										17.2	13.7	12.7	22.2		No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages

Linear Change[†]

Quadratic Change[†]

Change from
2017-2019[§]

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QNMILK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)

5.0 3.1 5.8 8.6 No linear change Not available[¶] No change

QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)

15.5 21.5 20.3 15.7 25.7 21.9 No linear change No quadratic change No change

QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)

23.5 25.7 27.9 28.5 22.1 18.2 No linear change No quadratic change No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Physical Activity**

Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]							
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															37.1	31.2	30.2	33.7	35.1	No linear change			Not available [¶]	No change
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															26.1	31.3	22.1	28.0	23.6	No linear change			Not available	No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															18.0	15.3	18.8	16.3	20.4	No linear change			Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Physical Activity

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																		
						59.6	44.9	45.1	46.1	38.0	32.6	29.9	29.3	Decreased, 2004-2019	No quadratic change	No change		
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																		
						29.5	26.1	33.1	37.8	30.0	34.4	43.4	Increased, 2007-2019	No quadratic change	No change			
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																		
						26.5	41.0	31.5	42.8	36.9	46.7	33.4	42.4	No linear change	No quadratic change	No change		

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Physical Activity

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)																	
						13.0	17.7	16.5	18.8	19.3	21.7	16.5	19.9		No linear change	No quadratic change	No change
QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)																	
						56.4	57.7	53.2	47.8	52.2	53.3	49.8	46.5		No linear change	No quadratic change	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Black* Other	Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017			
	QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)																
							57.4	57.2	64.1	66.1	48.3	61.4	56.8		No linear change	No quadratic change	No change
	QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)																
							5.6	3.1	5.7	4.8	5.9	3.5	4.8		No linear change	No quadratic change	No change
	QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma																
							26.5	35.1	26.8	37.3	30.5	33.2	30.2	37.4	No linear change	No quadratic change	No change
	QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)																
												16.2	18.7	14.1	No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black* Other	Health Risk Behavior and Percentages															Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]
	1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
	QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)																	
										58.0	66.2	54.8	70.9	67.8		No linear change	Not available [¶]	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Black*
Site-Added

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												52.4	54.7		No linear change	Not available [¶]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												45.5	49.5		No linear change	Not available	No change
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												39.5	36.0		No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[§]Based on t-test analysis, p < 0.05.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Site-Added**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]								
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)														23.4	13.4	17.5	No linear change	Not available [¶]	No change					
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)														5.9	4.0	4.7	5.8	3.6	15.4	6.4	5.2	No linear change	No quadratic change	No change
QN93: Percentage of students who had oral sex														41.8	56.0	41.1	36.6	36.7	34.5	No linear change	No quadratic change	No change		
QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)														15.1	11.8	No linear change	Not available	No change						

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Black*
Site-Added**

Health Risk Behavior and Percentages														Linear Change [†]	Quadratic Change [†]	Change from 2017-2019 [§]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																	
								70.9	79.0	72.0	70.6	61.5	64.7	66.3	Decreased, 2007-2019	No quadratic change	No change
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																	
								11.5	21.8	24.2	17.9	19.5	16.2	17.5	No linear change	No quadratic change	No change
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																	
													12.2	8.4	No linear change	Not available [¶]	No change
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																	
											25.0	15.4	10.0	16.3	No linear change	Not available	No change

*Non-Hispanic.

[†]Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[§]Based on t-test analysis, $p < 0.05$.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN8: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)																	
											17.6	20.1	15.2	10.2	No linear change	Not available [§]	No change
QN12: Percentage of students who carried a weapon (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
											18.1	26.0	14.9	14.8	No linear change	Not available	No change
QN13: Percentage of students who carried a weapon on school property (such as a gun, knife, or club, on at least 1 day during the 30 days before the survey)																	
											9.5	9.4	4.4	2.1	Decreased, 2013-2019	Not available	No change
QN14: Percentage of students who carried a gun (not counting the days when they carried a gun only for hunting or for a sport such as target shooting, on at least 1 day during the 12 months before the survey)																	
													10.9	8.3	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN16: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club, one or more times during the 12 months before the survey)														No linear change	Not available [§]	No change	
											10.4	9.5	9.2	10.9			
QN17: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)														No linear change	Not available	No change	
											28.2	23.8	24.1	25.7			
QN18: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)														No linear change	Not available	No change	
											11.9	9.4	11.8	6.7			

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN20: Percentage of students who experienced sexual violence (being forced by anyone to do sexual things [counting such things as kissing, touching, or being physically forced to have sexual intercourse] that they did not want to, one or more times during the 12 months before the survey)																	
												16.8	7.2	Decreased, 2017-2019	Not available [§]	Decreased	
QN23: Percentage of students who were bullied on school property (ever during the 12 months before the survey)																	
											26.2	20.4	17.2	22.5	No linear change	Not available	No change
QN24: Percentage of students who were electronically bullied (counting being bullied through texting, Instagram, Facebook, or other social media, ever during the 12 months before the survey)																	
											20.8	17.1	14.3	16.6	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN25: Percentage of students who felt sad or hopeless (almost every day for >=2 weeks in a row so that they stopped doing some usual activities, ever during the 12 months before the survey)																	
											31.5	31.4	30.0	38.6	No linear change	Not available [§]	No change
QN26: Percentage of students who seriously considered attempting suicide (ever during the 12 months before the survey)																	
											17.2	21.3	22.0	12.3	No linear change	Not available	No change
QN27: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)																	
											20.2	15.9	17.5	13.6	No linear change	Not available	No change
QN28: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)																	
											8.0	16.3	11.1	No linear change	Not available	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Injury and Violence

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN29: Percentage of students who had a suicide attempt that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)																	
												2.6	4.7	5.2	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN30: Percentage of students who ever tried cigarette smoking (even one or two puffs)																	
												48.6	45.7	28.1	Decreased, 2015-2019	Not available [§]	Decreased
QN31: Percentage of students who first tried cigarette smoking before age 13 years (even one or two puffs)																	
													15.5	9.0	No linear change	Not available	No change
QN32: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)																	
												25.6	12.4	6.9	Decreased, 2015-2019	Not available	No change
QNFRSIG: Percentage of students who currently smoked cigarettes frequently (on 20 or more days during the 30 days before the survey)																	
												12.5	1.6	2.3	Decreased, 2015-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]			
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019					
QNDAYCIG: Percentage of students who currently smoked cigarettes daily (on all 30 days during the 30 days before the survey)														9.2	1.2	1.3	No linear change	Not available [§]	No change
QN34: Percentage of students who ever used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo])														49.4	43.5	51.7	No linear change	Not available	No change
QN35: Percentage of students who currently used an electronic vapor product (including e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens [such as blu, NJOY, Vuse, MarkTen, Logic, Vapin Plus, eGo, and Halo], on at least 1 day during the 30 days before the survey)														25.6	16.0	23.8	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNFREVP: Percentage of students who currently used electronic vapor products frequently (on 20 or more days during the 30 days before the survey)																	
												5.9	4.9	4.0	No linear change	Not available [§]	No change
QNDAYEVP: Percentage of students who currently used electronic vapor products daily (on all 30 days during the 30 days before the survey)																	
												5.9	4.0	4.0	No linear change	Not available	No change
QNTB5: Percentage of students who currently smoked cigarettes or used electronic vapor products (on at least 1 day during the 30 days before the survey)																	
												36.2	19.2	26.1	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Tobacco Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, dip, snus, or dissolvable tobacco products [such as Copenhagen, Grizzly, Skoal, or Camel Snus], not counting any electronic vapor products, on at least 1 day during the 30 days before the survey)																	
												4.9	7.0		No linear change	Not available [§]	No change
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars, on at least 1 day during the 30 days before the survey)																	
										33.0	19.1	19.7	8.7	8.3	Decreased, 2011-2019	Not available	No change
QNTB2: Percentage of students who currently smoked cigarettes or cigars (on at least 1 day during the 30 days before the survey)																	
												29.8	14.1	12.3	Decreased, 2015-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN40: Percentage of students who had their first drink of alcohol before age 13 years (other than a few sips)																	
											25.8	25.5	20.6	20.1	No linear change	Not available [§]	No change
QN41: Percentage of students who currently drank alcohol (at least one drink of alcohol, on at least 1 day during the 30 days before the survey)																	
											33.4	26.1	24.2	No linear change	Not available	No change	
QN42: Percentage of students who currently were binge drinking (had four or more drinks of alcohol in a row for female students or five or more drinks of alcohol in a row for male students, within a couple of hours, on at least 1 day during the 30 days before the survey)																	
											10.2	8.7	No linear change	Not available	No change		

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN45: Percentage of students who ever used marijuana (one or more times during their life)														No linear change	Not available [§]	No change	
											39.0	41.5	36.9	41.0			
QN46: Percentage of students who tried marijuana for the first time before age 13 years														No linear change	Not available	No change	
											17.8	9.4	8.5	10.7			
QN47: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)														No linear change	Not available	No change	
											22.9	19.6	16.6	20.2			
QN48: Percentage of students who ever used synthetic marijuana (one or more times during their life)														No linear change	Not available	No change	
											18.1	10.7	13.7				

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN49: Percentage of students who ever took prescription pain medicine without a doctor's prescription or differently than how a doctor told them to use it (counting drugs such as codeine, Vicodin, OxyContin, Hydrocodone, and Percocet, one or more times during their life)																	
												12.5	13.8		No linear change	Not available [§]	No change
QN50: Percentage of students who ever used cocaine (any form of cocaine, including powder, crack, or freebase, one or more times during their life)																	
											13.4	10.0	8.1	6.2	No linear change	Not available	No change
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)																	
											13.2	16.0	8.2	5.6	Decreased, 2013-2019	Not available	No change
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China White," one or more times during their life)																	
												8.7	3.6	3.7	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic Alcohol and Other Drug Use

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)																	
											11.3	15.0	6.2	4.4	Decreased, 2013-2019	Not available [§]	No change
QN55: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)																	
											10.5	10.6	3.5	3.0	Decreased, 2013-2019	Not available	No change
QN56: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body, one or more times during their life)																	
													3.7	5.7	No linear change	Not available	No change
QN57: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)																	
											33.2	23.7	28.5	29.6	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN58: Percentage of students who ever had sexual intercourse														No linear change	Not available [§]	No change	
												43.0	39.4	47.8			
QN59: Percentage of students who had sexual intercourse for the first time before age 13 years														No linear change	Not available	No change	
												14.3	7.0	7.0			
QN60: Percentage of students who had sexual intercourse with four or more persons during their life														No linear change	Not available	No change	
												19.0	9.8	9.8			
QN61: Percentage of students who were currently sexually active (had sexual intercourse with at least one person, during the 3 months before the survey)														No linear change	Not available	No change	
												30.9	25.6	37.0			

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Sexual Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN84: Percentage of students who were ever tested for human immunodeficiency virus (HIV) (not counting tests done if they donated blood)																	
												22.4	13.5	11.1	No linear change	Not available [§]	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019											
QNOWT: Percentage of students who were overweight (\geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]														17.0	11.8	18.6	17.2					No linear change	Not available [¶]	No change	
QNOBESE: Percentage of students who had obesity (\geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts) [§]														18.8	19.0	17.7	14.3					No linear change	Not available	No change	
QN67: Percentage of students who described themselves as slightly or very overweight														40.3	28.4	27.2	35.7	31.5					No linear change	Not available	No change
QN69: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)														21.7	18.9	30.2	33.1	32.3					Increased, 2011-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Overweight and obese prevalence estimates for 1999 differ slightly from previously published results because different BMI cut points were used in 1999 than in subsequent years. To make these prevalence estimates comparable, the 1999 prevalence estimates were recalculated using the updated BMI cut points. In addition, beginning in 2017, new, slightly different ranges were used to calculate biologically implausible responses to height and weight questions.

[¶]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN70: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)																	
										21.6	9.8	15.8	14.5	13.2	No linear change	Not available [§]	No change
QNFR0: Percentage of students who did not eat fruit or drink 100% fruit juices (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
										11.5	7.1	12.3	11.7	9.0	No linear change	Not available	No change
QNFR1: Percentage of students who ate fruit or drank 100% fruit juices one or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)																	
										60.1	73.2	47.9	47.1	52.3	Decreased, 2011-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]										
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019													
QNFR2: Percentage of students who ate fruit or drank 100% fruit juices two or more times per day (such as orange juice, apple juice, or grape juice, during the 7 days before the survey)															35.5	43.7	26.5	32.0	30.4						No linear change	Not available [§]	No change
QN71: Percentage of students who did not eat green salad (one or more times during the 7 days before the survey)															46.7	42.6	44.7	52.2	49.7						No linear change	Not available	No change
QN72: Percentage of students who did not eat potatoes (one or more times during the 7 days before the survey)															21.1	27.8	37.1	43.4	40.0						Increased, 2011-2019	Not available	No change
QN73: Percentage of students who did not eat carrots (one or more times during the 7 days before the survey)															59.2	60.8	53.7	60.4						No linear change	Not available	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN74: Percentage of students who did not eat other vegetables (one or more times during the 7 days before the survey)																	
								25.3	24.1	29.0	30.8	23.9			No linear change	Not available [§]	No change
QNVEG0: Percentage of students who did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
											8.9	10.9	15.9	12.2	No linear change	Not available	No change
QNVEG1: Percentage of students who ate vegetables one or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)																	
											54.4	50.9	49.4	52.8	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †									
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019											
QNVEG2: Percentage of students who ate vegetables two or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)														20.4	24.7	23.1	23.7					No linear change	Not available [§]	No change	
QNVEG3: Percentage of students who ate vegetables three or more times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)														10.0	14.9	14.3	13.3					No linear change	Not available	No change	
QN75: Percentage of students who did not drink a can, bottle, or glass of soda or pop (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)														15.6	29.7	27.2	30.0	31.2					Increased, 2011-2019	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

†Based on t-test analysis, $p < 0.05$.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]								
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019										
QNSODA1: Percentage of students who drank a can, bottle, or glass of soda or pop one or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)														33.8	19.6	24.1	21.1	17.6				No linear change	Not available [§]	No change
QNSODA2: Percentage of students who drank a can, bottle, or glass of soda or pop two or more times per day (such as Coke, Pepsi, or Sprite, not counting diet soda or diet pop, during the 7 days before the survey)														29.3	14.2	16.8	15.3	13.7				No linear change	Not available	No change
QN76: Percentage of students who did not drink milk (during the 7 days before the survey)														20.6	20.1	23.9	28.4				No linear change	Not available	No change	
QNMILK1: Percentage of students who drank one or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)														29.1	26.4	27.6	23.9				No linear change	Not available	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

[†]Based on t-test analysis, $p < 0.05$.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic

Weight Management and Dietary Behaviors

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNMILK3: Percentage of students who drank three or more glasses per day of milk (counting the milk they drank in a glass or cup, from a carton, or with cereal and counting the half pint of milk served at school as equal to one glass, during the 7 days before the survey)														No linear change	Not available [§]	No change	
											12.0	9.2	7.8	7.4			
QN77: Percentage of students who did not eat breakfast (during the 7 days before the survey)														No linear change	Not available	No change	
											15.5	14.6	18.0	17.9			
QNBK7DAY: Percentage of students who ate breakfast on all 7 days (during the 7 days before the survey)														No linear change	Not available	No change	
											35.4	33.5	31.1	29.4			

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Physical Activity

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN78: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															No linear change	Not available [§]	No change
QNPA0DAY: Percentage of students who did not participate in at least 60 minutes of physical activity on at least 1 day (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															No linear change	Not available	No change
QNPA7DAY: Percentage of students who were physically active at least 60 minutes per day on all 7 days (in any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)															No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Physical Activity

Health Risk Behavior and Percentages															Linear Change*	Quadratic Change*	Change from 2017-2019 [†]
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN79: Percentage of students who watched television 3 or more hours per day (on an average school day)																	
										39.1	21.6	32.7	27.1	20.8	No linear change	Not available [§]	No change
QN80: Percentage of students who played video or computer games or used a computer 3 or more hours per day (counting time spent on things such as Xbox, PlayStation, an iPad or other tablet, a smartphone, texting, YouTube, Instagram, Facebook, or other social media, for something that was not school work, on an average school day)																	
											29.7	43.6	45.1	41.9	No linear change	Not available	No change
QN81: Percentage of students who attended physical education (PE) classes on 1 or more days (in an average week when they were in school)																	
										48.7	39.4	37.6	40.6	35.4	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Physical Activity

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019†	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNDLYPE: Percentage of students who attended physical education (PE) classes on all 5 days (in an average week when they were in school)														No linear change	Not available§	No change	
QN82: Percentage of students who played on at least one sports team (counting any teams run by their school or community groups, during the 12 months before the survey)														Decreased, 2011-2019	Not available	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Other

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work, during the 12 months before the survey)														No linear change	Not available [§]	No change	
											63.5	65.8	65.8	63.2			
QNNODNT: Percentage of students who never saw a dentist (for a check-up, exam, teeth cleaning, or other dental work)														No linear change	Not available	No change	
											4.5	6.4	3.8	4.3			
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma														No linear change	Not available	No change	
											22.9	18.3	15.1	22.4			
QN88: Percentage of students who got 8 or more hours of sleep (on an average school night)														No linear change	Not available	No change	
											21.3	17.6	22.3				

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Other

Health Risk Behavior and Percentages

Linear Change*

Quadratic Change*

Change from
2017-2019 †

1991 1993 1995 1997 1999 2001 2003 2004 2007 2009 2011 2013 2015 2017 2019

QN89: Percentage of students who described their grades in school as mostly A's or B's (during the 12 months before the survey)

61.3 57.5 64.9 70.5 No linear change Not available‡ No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, $p < 0.05$.

†Based on t-test analysis, $p < 0.05$.

‡Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Site-Added

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 †	
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019			
QNWATER1: Percentage of students who drank a bottle or glass of plain water one or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												55.7	66.0		No linear change	Not available [§]	No change
QNWATER2: Percentage of students who drank a bottle or glass of plain water two or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												51.8	58.6		No linear change	Not available	No change
QNWATER3: Percentage of students who drank a bottle or glass of plain water three or more times per day (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																	
												40.5	46.0		No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

†Based on t-test analysis, p < 0.05.

§Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey Trend Analysis Report

**Hispanic
Site-Added**

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]		
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019				
QN90: Percentage of students who did something to purposely hurt themselves without wanting to die (such as cutting or burning themselves on purpose one or more times during the 12 months before the survey)																		
											26.4	22.6	17.9		No linear change	Not available [§]	No change	
QN91: Percentage of students who smoked cigarettes on school property (on at least 1 day during the 30 days before the survey)																		
											6.4	14.0	6.6	5.7		No linear change	Not available	No change
QN93: Percentage of students who had oral sex																		
											49.2	41.7	35.8	41.4		No linear change	Not available	No change
QN94: Percentage of students who did not drink a bottle or glass of plain water (counting tap, bottled, and unflavored sparkling water, during the 7 days before the survey)																		
												16.6	9.6		No linear change	Not available	No change	

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.

2019 YOUTH RISK BEHAVIOR SURVEY RESULTS

Kentucky High School Survey
Trend Analysis Report

Hispanic
Site-Added

Health Risk Behavior and Percentages														Linear Change*	Quadratic Change*	Change from 2017-2019 [†]															
1991	1993	1995	1997	1999	2001	2003	2004	2007	2009	2011	2013	2015	2017	2019																	
QN95: Percentage of students who brushed their teeth on all 7 days (during the 7 days before the survey)																															
															72.5	62.9	68.2	69.5	No linear change	Not available [§]	No change										
QN96: Percentage of students who flossed their teeth on all 7 days (during the 7 days before the survey)																															
															19.5	22.1	17.6	20.9	No linear change	Not available	No change										
QN99: Percentage of students who did not usually sleep in their parent's or guardian's home (during the 30 days before the survey)																															
																						11.2	7.7	No linear change	Not available	No change					
QN100: Percentage of students who do not have an adult in their life that they trust and can talk to about serious problems																															
																										16.7	13.0	19.5	No linear change	Not available	No change

*Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade, p < 0.05.

[†]Based on t-test analysis, p < 0.05.

[§]Not enough years of data to calculate.